



On Two Absolute Index-Summability Methods

B.P.Padhy, Dattaram Bisoyi, Mahendra Misra, U.K.Misra

Department of Mathematics, Roland Institute of Technology,Golanthara-761008, Odisha, India

Email:iraady@gmail.com

Department of Mathematics, L.N.Mahavidyalaya,Kodala,Ganjam, Odisha,India

Email: dbisoyi2@gmail.com

Department of Mathematics,Binayak Acharya College,Berhampur, Odisha, India

Email: mahendramisra@2007.gmail.com

Department of Mathematics, National Institute of Science and Technology,PallurHills-761008, Odisha, India

Email: umakanta_misra@yahoo.com

ABSTRACT: In this paper we have established a relation between the Summability methods $X - \left| \bar{N}, p_n \right|_k$,

$Y - |A|_k$, $k \geq 1$ and $Y - |A, f(\delta)|_k$, $k \geq 1, \delta \geq 0$.

Keywords: $\left| \bar{N}, p_n \right|_k$, $X - \left| \bar{N}, p_n \right|_k$, $X - |N, p_n|_k$, $X - |A|_k$ -summabilities.

AMS Subject Classification: 40D25



Council for Innovative Research

Peer Review Research Publishing System

Journal: Journal of Advances in Mathematics

Vol 6, No. 1

editor@cirworld.com

www.cirworld.com, member.cirworld.com



1. Introduction:

Let $\sum a_n$ be an infinite series and $\{s_n\}$ the sequence of partial sums. Let $\{p_n\}$ be a sequence of non-negative numbers with $P_n = \sum_{v=0}^n p_v$ for all $n \in N$. The sequence-to-sequence transformation

$$(1.1) \quad t_n = \frac{1}{P_n} \sum_{v=0}^n p_v s_v, P_n \neq 0$$

defines $|N, p_n|$ -mean of the sequence $\{s_n\}$ generated by the sequence of coefficients $\{p_n\}$. The series $\sum a_n$ is said to be summable $|N, p_n|_k$, $k \geq 1$, [4] if

$$(1.2) \quad \sum_{n=1}^{\infty} \left(\frac{P_n}{p_n} \right)^{k-1} |t_n - t_{n-1}|^k < \infty,$$

The sequence-to-sequence transformation

$$(1.3) \quad \tau_n = \frac{1}{P_n} \sum_{v=0}^n p_{n-v} s_v, P_n \neq 0,$$

defines $|N, p_n|$ -mean of the sequence $\{s_n\}$. The series $\sum a_n$ is said to be summable $|N, p_n|_k$, $k \geq 1$, if

$$(1.4) \quad \sum_{n=1}^{\infty} \left(\frac{P_n}{p_n} \right)^{k-1} |\tau_n - \tau_{n-1}|^k < \infty.$$

The series $\sum a_n$ is said to be summable $X - |N, p_n|_k$, $k \geq 1$, if

$$(1.5) \quad \sum_{n=1}^{\infty} X_n^{k-1} |t_n - t_{n-1}|^k < \infty$$

where $\{X_n\}$ is a sequence of positive real constants. Similarly, $\sum a_n$ is said to be summable $X - |N, p_n|_k$, $k \geq 1$, if

$$(1.6) \quad \sum_{n=1}^{\infty} X_n^{k-1} |\tau_n - \tau_{n-1}|^k < \infty.$$

Let $A = (a_{nk})$ be a $\infty \times \infty$ matrix. The series $\sum a_n$ is said to be summable $X - |A|_k$, $k \geq 1$, if

$$(1.7) \quad \sum_{n=1}^{\infty} X_n^{k-1} |T_n - T_{n-1}|^k < \infty,$$

$\sum a_n$ is said to be summable $X - |A, \delta|_k$, $k \geq 1, \delta \geq 0$, if

$$(1.8) \quad \sum_{n=1}^{\infty} X_n^{k+k-1} |T_n - T_{n-1}|^k < \infty$$

and $\sum a_n$ is said to be summable $X - |A, f(\delta)|_k$, $k \geq 1, \delta \geq 0$, if



$$(1.9) \quad \sum_{n=1}^{\infty} (f(\delta))^k X_n^{k-1} |T_n - T_{n-1}|^k < \infty,$$

where the sequence-to-sequence transformation $\{T_n\}$ is given by

$$(1.10) \quad T_n = \sum_{k=1}^{\infty} a_{nk} s_k.$$

Clearly $X - |A, f(\delta)|_k, k \geq 1, \delta \geq 0$ reduces to $X - |A, \delta|_k, k \geq 1, \delta \geq 0$ if $f(\delta) = X_n^\delta$.

2. Known Theorems:

Dealing with the index summability method Bor has established the following theorems:

Theorem-A[1]:

Let $\{p_n\}$ be a sequence of positive real constants such that as $n \rightarrow \infty$

$$\text{i)} np_n = O(P_n) \quad \text{ii)} P_n = O(np_n).$$

If $\sum a_n$ is summable $|C, 1|_k$ then it is summable $|\bar{N}, p_n|_k, k \geq 1$.

Theorem-B[2]:

Let $\{p_n\}$ be a sequence of positive real constants such that as $n \rightarrow \infty$

$$\text{i)} np_n = O(P_n) \quad \text{ii)} P_n = O(np_n).$$

If $\sum a_n$ is summable $|\bar{N}, p_n|_k$ then it is summable $|C, 1|_k, k \geq 1$.

Subsequently Bor and Thorpe established the following result.

Theorem-C[3]:

Let $\{p_n\}$ and $\{q_n\}$ be the sequence of positive real constants such that

$$\text{i)} p_n Q_n = O(P_n q_n) \quad \text{ii)} P_n q_n = O(p_n Q_n).$$

then the series $\sum a_n$ is summable $|\bar{N}, q_n|_k$ whenever it is summable $|\bar{N}, p_n|_k, k \geq 1$.

Further, Tripathi established

Theorem-D[6]:

Suppose $\{p_n\}, \{q_n\}, \{X_n\}$ and $\{Y_n\}$ are sequences of positive real constants such that

$$\text{i)} q_n p_n = O(p_n q_n) \quad \text{ii)} Q_n = O(q_n X_n) \quad \text{iii)} Y_n p_n = O(p_n).$$

If $\sum a_n$ is summable $X - |\bar{N}, p_n|_k$, then it is summable $Y - |\bar{N}, q_n|_k, k \geq 1$.

Extending the above result, Misra, Misra and Routa established the following theorem replacing $Y - |\bar{N}, q_n|_k, k \geq 1$ by $Y - |N, q_n|_k, k \geq 1$, in Theorem-D.

Theorem-E[7]:

Suppose $\{p_n\}, \{q_n\}, \{X_n\}$ and $\{Y_n\}$ are sequences of positive real constants such that

$$\text{i)} q_n p_n = O(p_n q_n) \quad \text{ii)} Q_n = O(q_n X_n) \quad \text{iii)} Y_n p_n = O(p_n)$$



$$\text{iv)} \sum_{n=v+1}^{n+1} \frac{1}{Q_n} = O\left(\frac{1}{Q_\gamma}\right) \quad \text{v)} \sum_{v=1}^{n-1} a_v^{\frac{k}{k-1}} = O(1), k \neq 1.$$

If $\sum a_n$ is summable $X - |\bar{N}, p_n|_k$, then it is summable $Y - |\bar{N}, q_n|_k$, $k \geq 1$.

Further, generalizing the above theorem to matrix summability Padhy, panda, Misra and Misra established the following theorem:

Theorem-F[5]:

Suppose $\{p_n\}$, $\{q_n\}$, $\{X_n\}$ and $\{Y_n\}$ are sequences of positive real constants such that

$$\text{i)} \quad a_{nn}P_n = O(p_n), \quad \text{ii)} \quad \frac{1}{a_{nn}} = O(X_n), \quad \text{iii)} \quad Y_n p_n = O(P_n),$$

$$\text{iv)} \sum_{n=r}^{m+1} \left(\frac{P_n}{P_r} \right)^{\delta k + k - 1} A_{nr} = O(a_{rr}) \quad , \text{ where } A_{nk} = \sum_{v=k}^n a_{nv}$$

and, for the infinite triangular matrix $A = (a_{nk})_{\infty \times \infty}$,

$$\text{v)} \sum_{r=1}^n A_{nr} = O(1).$$

Then $\sum a_n$ is $Y - |A, \delta|_k$, $k \geq 1, \delta \geq 0$ summable whenever $\sum a_n$ is summable $X - |\bar{N}, p_n|_k$, $k \geq 1$.

In what follows, in the present paper, further generalizing the above theorem we establish

3. Main Theorem:

Suppose $\{p_n\}$, $\{q_n\}$, $\{X_n\}$ and $\{Y_n\}$ are sequences of positive real constants such that

$$(3.1) \quad \text{i)} \quad a_{nn}P_n = O(p_n),$$

$$(3.2) \quad \text{ii)} \quad \frac{1}{a_{nn}} = O(X_n),$$

$$(3.3) \quad \text{iii)} \quad Y_n p_n = O(P_n),$$

$$(3.4) \quad \text{iv)} \sum_{n=r}^{m+1} (f(\delta))^k \left(\frac{P_n}{P_r} \right)^{k-1} A_{nr} = O(a_{rr}) \quad , \text{ where } A_{nk} = \sum_{v=k}^n a_{nv}$$

and, for the infinite triangular matrix $A = (a_{nk})_{\infty \times \infty}$,

$$(3.5) \quad \text{v)} \sum_{r=1}^n A_{nr} = O(1).$$

Then $\sum a_n$ is $Y - |A, f(\delta)|_k$, $k \geq 1, \delta \geq 0$ summable, whenever $\sum a_n$ is summable $X - |\bar{N}, p_n|_k$, $k \geq 1$.

4. Proof of Theorem:

If $\{t_n\}$ is the nth $|\bar{N}, p_n|$ -mean of $\sum a_n$, then



$$\begin{aligned}
t_n &= \frac{1}{P_n} \sum_{\nu=0}^n p_\nu s_\nu \\
&= \frac{1}{P_n} \{p_0 a_0 + p_1 (a_0 + a_1) + \cdots + p_n (a_0 + a_1 + \cdots + a_n)\} \\
&= \frac{1}{P_n} \{P_n a_0 + (P_n - p_0) a_1 + \cdots + (P_n - p_{n-1}) a_n\} \\
&= \frac{1}{P_n} \sum_{\nu=0}^n (P_n - P_{\nu-1}) a_\nu
\end{aligned}$$

Then

$$\begin{aligned}
\Delta t_n &= t_n - t_{n-1} \\
&= \frac{1}{P_n} \sum_{\nu=0}^n (P_n - P_{\nu-1}) a_\nu - \frac{1}{P_{n-1}} \sum_{\nu=0}^{n-1} (P_{n-1} - P_{\nu-1}) a_\nu \\
&= \frac{1}{P_n} \sum_{\nu=1}^n (P_n - P_{\nu-1}) a_\nu - \frac{1}{P_{n-1}} \sum_{\nu=0}^{n-1} (P_{n-1} - P_{\nu-1}) a_\nu \\
&= \left(\frac{1}{P_{n-1}} - \frac{1}{P_n} \right) \sum_{\nu=1}^n P_{\nu-1} a_\nu \\
&= \frac{P_n}{P_n P_{n-1}} \sum_{\nu=1}^n P_{\nu-1} a_\nu
\end{aligned}$$

Hence,

$$\frac{P_n P_{n-1}}{p_n} \Delta t_n = \sum_{\nu=1}^n P_{\nu-1} a_\nu$$

and

$$\frac{P_{n-1} P_{n-1}}{p_{n-1}} \Delta t_{n-1} = \sum_{\nu=1}^{n-1} P_{\nu-1} a_\nu$$

Thus,

$$a_n = \frac{P_n}{p_n} \Delta t_n - \frac{P_{n-2}}{p_{n-1}} \Delta t_{n-1}$$

Further, if $\{T_n\}$ is the nth A-mean of $\sum a_n$, where $A = (a_{nk})_{\infty \times \infty}$ triangular matrix,

$$\begin{aligned}
T_n &= \sum_{k=0}^n a_{nk} s_k \\
&= \sum_{k=0}^n A_{nk} a_k, \quad A_{nk} = \sum_{\nu=k}^n a_{n\nu}
\end{aligned}$$

Then,



$$\begin{aligned}
T_n - T_{n-1} &= \sum_{k=0}^n A_{nk} a_k - \sum_{k=0}^{n-1} A_{n-1,k} a_k \\
&= \sum_{k=1}^n (A_{nk} - A_{n-1,k}) a_k \\
&= \sum_{k=1}^n (A_{nk} - A_{n-1,k}) \left(\frac{P_k}{p_k} \Delta t_k - \frac{P_{k-2}}{p_{k-1}} \Delta t_{k-1} \right) \\
&= \sum_{k=1}^n A_{nk} \frac{P_k}{p_k} \Delta t_k - \sum_{k=1}^n A_{nk} \frac{P_{k-2}}{p_{k-1}} \Delta t_{k-1} - \sum_{k=1}^{n-1} A_{n-1,k} \frac{P_k}{p_k} \Delta t_k + \sum_{k=1}^{n-1} A_{n-1,k} \frac{P_{k-2}}{p_{k-1}} \Delta t_{k-1} \\
&= S_1 + S_2 + S_3 + S_4 \text{ (say).}
\end{aligned}$$

Now,

$$\sum_{n=1}^{m+1} (f(\delta))^k Y_n^{k-1} |T_n - T_{n-1}|^k \leq \sum_{n=1}^{m+1} (f(\delta))^k Y_n^{k-1} |S_1 + S_2 + S_3 + S_4|^k = \sum_{i=1}^4 \sum_{n=1}^{m+1} Y_n^{\delta k+k-1} |S_i|^k$$

(By Minokowski's inequality)

Our Theorem will be established if we show that $\sum_{n=1}^{m+1} Y_n^{\delta k+k-1} |S_i|^k < \infty$, $\forall i = 1, 2, 3, 4$.

$$\begin{aligned}
\sum_{n=1}^{m+1} (f(\delta))^k Y_n^{k-1} |S_1|^k &= \sum_{n=1}^{m+1} (f(\delta))^k Y_n^{k-1} \left| \sum_{r=1}^n A_{nr} \frac{P_r}{p_r} \Delta t_r \right|^k \\
&\leq \sum_{n=1}^{m+1} (f(\delta))^k Y_n^{k-1} \sum_{r=1}^n \left(\frac{P_r}{p_r} \right)^k |\Delta t_r|^k A_{nr} \left(\sum_{r=1}^n A_{nr} \right)^{k-1}
\end{aligned}$$

(Using Holder's inequality)

$$\begin{aligned}
&= O(1) \sum_{r=1}^{m+1} \left(\frac{P_r}{p_r} \right)^k |\Delta t_r|^k \sum_{n=r}^{m+1} (f(\delta))^k Y_n^{k-1} A_{nr}, \text{ by (3.5)} \\
&= O(1) \sum_{r=1}^{m+1} \left(\frac{P_r}{p_r} \right)^k |\Delta t_r|^k \sum_{n=r}^{m+1} (f(\delta))^k \left(\frac{P_n}{p_n} \right)^{k-1} A_{nr}, \text{ by (3.3)} \\
&= O(1) \sum_{r=1}^{m+1} \left(\frac{1}{a_{rr}} \right)^k |\Delta t_r|^k a_{rr}, \text{ (using 3.4)} \\
&= O(1) \sum_{r=1}^{m+1} X_r^{k-1} |\Delta t_r|^k, \text{ (using 3.2)} \\
&= O(1).
\end{aligned}$$

Next

$$\sum_{n=1}^{m+1} (f(\delta))^k Y_n^{k-1} |S_2|^k = \sum_{n=1}^{m+1} (f(\delta))^k Y_n^{k-1} \left| \sum_{r=1}^n A_{nr} \frac{P_{r-2}}{p_{r-1}} \Delta t_{r-1} \right|^k$$



$$\begin{aligned}
&\leq \sum_{n=1}^{m+1} (f(\delta))^k Y_n^{k-1} \sum_{r=1}^n \left(\frac{P_{r-1}}{p_{r-1}} \right)^k |\Delta t_{r-1}|^k A_{nr} \left(\sum_{r=1}^n A_{nr} \right)^{k-1} \\
&= O(1) \sum_{r=1}^{m+1} \left(\frac{P_{r-1}}{p_{r-1}} \right)^k |\Delta t_{r-1}|^k \sum_{n=r}^{m+1} (f(\delta))^k Y_n^{k-1} A_{nr}, \text{ by (3.5)} \\
&= O(1) \sum_{r=1}^{m+1} \left(\frac{P_r}{p_{r-1}} \right)^k |\Delta t_{r-1}|^k \sum_{n=r}^{m+1} (f(\delta))^k \left(\frac{P_n}{p_n} \right)^{k-1} A_{nr}, \text{ by (3.3)} \\
&= O(1) \sum_{r=2}^{m+1} \left(\frac{P_r}{p_r} \right)^k |\Delta t_r|^k a_{rr}, \text{ (using 3.4)} \\
&= O(1) \sum_{r=2}^{m+1} \left(\frac{1}{a_{rr}} \right)^k |\Delta t_r|^k a_{rr} \\
&= O(1) \sum_{r=2}^{m+1} X_r^{k-1} |\Delta t_r|^k, \text{ (using 3.2)} \\
&= O(1).
\end{aligned}$$

Also,

$$\begin{aligned}
\sum_{n=1}^{m+1} (f(\delta))^k Y_n^{k-1} |S_3|^k &= \sum_{n=1}^{m+1} (f(\delta))^k Y_n^{k-1} \left| \sum_{r=1}^{n-1} A_{n-1,r} \frac{P_r}{p_r} \Delta t_r \right|^k \\
&\leq \sum_{n=1}^{m+1} (f(\delta))^k Y_n^{k-1} \sum_{r=1}^{n-1} \left(\frac{P_r}{p_r} \right)^k |\Delta t_r|^k A_{n-1,r} \left(\sum_{r=1}^n A_{n-1,r} \right)^{k-1} \\
&= O(1) \sum_{r=1}^{m+1} \left(\frac{P_r}{p_r} \right)^k |\Delta t_r|^k \sum_{n=r+1}^{m+1} (f(\delta))^k \left(\frac{P_n}{p_n} \right)^{k-1} A_{n-1,r}, \text{ using (3.5)} \\
&= O(1) \sum_{r=1}^{m+1} \left(\frac{1}{a_{rr}} \right)^k |\Delta t_r|^k a_{rr}, \text{ (using 3.4)} \\
&= O(1) \sum_{r=1}^{m+1} X_r^{k-1} |\Delta t_r|^k, \text{ (using 3.2)} \\
&= O(1).
\end{aligned}$$

Finally,

$$\begin{aligned}
\sum_{n=1}^{m+1} (f(\delta))^k Y_n^{k-1} |S_4|^k &= \sum_{n=1}^{m+1} (f(\delta))^k Y_n^{k-1} \left| \sum_{r=1}^{n-1} A_{n-1,r} \frac{P_{r-2}}{p_{r-2}} \Delta t_{r-1} \right|^k \\
&\leq \sum_{n=1}^{m+1} (f(\delta))^k Y_n^{k-1} \sum_{r=1}^{n-1} \left(\frac{P_{r-2}}{p_{r-2}} \right)^k |\Delta t_{r-1}|^k A_{n-1,r} \left(\sum_{r=1}^n A_{n-1,r} \right)^{k-1}
\end{aligned}$$



$$\begin{aligned} &= O(1) \sum_{n=1}^{m+1} (f(\delta))^k Y_n^{k-1} \sum_{r=1}^{n-1} \left(\frac{P_{r-1}}{p_{r-1}} \right)^k |\Delta t_{r-1}|^k A_{n-1,r} \\ &= O(1) \sum_{r=1}^m \left(\frac{P_{r-1}}{p_{r-1}} \right)^k |\Delta t_{r-1}|^k \sum_{n=r+1}^{m+1} A_{n-1,r} (f(\delta))^k Y_n^{k-1} \\ &= O(1) \sum_{r=1}^m \left(\frac{1}{a_{r-1,r-1}} \right)^k |\Delta t_{r-1}|^k a_{r-1,r-1}, \text{ (using 3.4)} \\ &= O(1) \sum_{r=1}^m X_{r-1}^{k-1} |\Delta t_{r-1}|^k, \text{ (using 3.2)} \\ &= O(1). \end{aligned}$$

References

- [1] Hussain Bor, "A note On two Summability methods", Math.Proc. Camb. Phil.Soc.,Vol.97(1985),147-149.
- [2] Hussain Bor, "A note On two Summability methods", Proc.Amer. Math. Soc.,Vol.98(1986),81-84.
- [3] Hussain Bor and Thorpe, "On some Summability methods" ,Analysis, Vol.7(1987),Pp-147-152.
- [4] S.M .Mazher, "Absolute Summability factor for Fourier Series" ,Indian J.Math.,Vol.41,No.2(1999),221-229.
- [5] B.P.Padhy, N.K.Panda, U.K.Misra and M.Misra, "On Relation between two Absolute index-Summability Methods", International Mathematical Forum, Vol.7, No.60(2012), 2977 – 2986.
- [6] Sanjaya Tripathy, "A Relation between two Summability Methods", Bulletin of Pure and Applied Sciences.Vol.20E, No.1 (2001).
- [7] U.K .Misra, M.Misra, K.Rauto"Relation between two Summability Methods" ,Acta Ciencia Indica,Vol.XXI M, No.4 (2005), 1173-1179.