

On Twisted Sums of Sequence Spaces.

Haifa Bin Jebreen

Mathematics department, College of Science, King Saud University.
hjebreen@ksu.edu.sa

Abstract

We prove the existence of non trivial twisted sums involving the p^{th} James space J_p ($1 \leq p < \infty$), the Johnson-Lindenstrauss space JL , the James tree space JT , the Tsirelson's space T and the Argyros and Deliyanni space AD . We also present non trivial twisted sums involving their duals and biduals. We show that there are strictly singular quasi-linear maps from the spaces T, T^*, AD and JT into $C[0, 1]$. We discuss the Pelczynski's property (u) for the twisted sums involving these spaces which extends a p^{th} James-Schreier spaces V_p ($1 < p < \infty$) or J_2 .

Keywords: Twisted sums, James spaces, Tsirlson's spaces, strictly singular.

Subject Classification: 46B03; 46B20; 46B45.

Introduction

A quasi-Banach space X is said to be a twisted sum of two Banach spaces Y and U if it contains a subspace A isomorphic to Y and the quotient X/A is isomorphic to U . Identifying A with Y we have the following short exact sequence

$$0 \longrightarrow Y \longrightarrow X \longrightarrow U \longrightarrow 0$$

Two exact sequences $0 \longrightarrow Y \longrightarrow X_1 \longrightarrow U \longrightarrow 0$ and $0 \longrightarrow Y \longrightarrow X_2 \longrightarrow U \longrightarrow 0$ are equivalent if there is a bounded linear operator T making the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & Y & \longrightarrow & X_1 & \longrightarrow & U \longrightarrow 0 \\ & & & & \parallel & T \downarrow & \parallel \\ 0 & \longrightarrow & Y & \longrightarrow & X_2 & \longrightarrow & U \longrightarrow 0 \end{array}$$

commutative. The three-lemma and the open mapping theorem imply that T must be an isomorphism [10, 1.5]. An exact sequence $0 \longrightarrow Y \longrightarrow X \longrightarrow U \longrightarrow 0$ is said to split and X is said to be trivial if it is equivalent to the trivial sequence $0 \longrightarrow Y \longrightarrow Y \oplus U \longrightarrow U \longrightarrow 0$.

A quasi-linear map $F : U \rightarrow Y$ where U and Y are Banach spaces is a homogeneous map such that

$$\|F(u+z) - F(u) - F(z)\| \leq k(\|u\| + \|z\|)$$



for some constant k and all $u, z \in U$. For a quasi-linear map $F : U \rightarrow Y$, there corresponds a twisted sum $Y \oplus_F U$ by endowing the product space $Y \times U$ with the quasi-norm $\|(y, z)\| = \|y - F(z)\| + \|z\|$. The subspace $\{(y, 0) : y \in Y\}$ of $Y \oplus_F U$ is isometric to Y and the corresponding quotient $(Y \oplus_F U)/Y$ is isomorphic to U . Conversely, for every twisted sum of Y and U there is a quasi-linear map $F : U \rightarrow Y$ such that X is equivalent to $Y \oplus_F U$ [10, 1.5]. Two quasi-linear maps F and G of a Banach space U into a Banach space Y are said to be equivalent if the corresponding twisted sums $Y \oplus_F U$ and $Y \oplus_G U$ are equivalent. If the quasi-linear map $F : U \rightarrow Y$, acting between two Banach spaces U and Y , is zero-linear, that is F satisfies

$$\left\| F\left(\sum_{i=1}^n u_i\right) - \sum_{i=1}^n F(u_i) \right\| \leq k \left(\sum_{i=1}^n \|u_i\|\right).$$

for some constant k , where u_1, u_2, \dots, u_n are finitely many elements of U , then the twisted sum $Y \oplus_F U$ is locally convex [10, 1.6.e]. We denote by $Ext(U, Y)$ the space of all equivalence classes of locally convex twisted sums of Y and U . Thus $Ext(U, Y) = 0$ means that all locally convex twisted sums of Y and U are equivalent to the direct sum $Y \oplus U$.

Given a family \mathcal{E} of finite dimensional Banach spaces, a Banach space X is said to contain \mathcal{E} uniformly complemented if there exists a constant c such that for every $E \in \mathcal{E}$, there is a c -complemented subspace A of X which is c -isomorphic to E . It is clear that X contains \mathcal{E} uniformly complemented if and only if its second dual X^{**} does. A Banach space X is said to be λ -locally \mathcal{E} (or locally \mathcal{E}) if there exists a constant $\lambda > 1$ such that every finite dimensional subspace A of X is contained in a finite dimensional subspace B of X such that

$$d_{BM}(B, E) = \inf\{\|T\| \|T^{-1}\|; T : X \rightarrow Y \text{ is an isomorphism of } X \text{ onto } Y\} < \lambda$$

for some $E \in \mathcal{E}$ [6].

We say that a Banach space X is λ -colocally \mathcal{E} (or colocally \mathcal{E}) if there exists a constant $\lambda > 1$ such that every finite dimensional quotient A of X is a quotient of another finite dimensional quotient B of X satisfying $d_{BM}(B, E) < \lambda$ for some $E \in \mathcal{E}$ [17].

The locality of a family is a very useful tool to determine the existence of nontrivial twisted sums of certain Banach spaces, in fact, Cabello and Castillo proved

Theorem 1 [6, Theorem 2] *Let \mathcal{E} be a family of finite dimensional Banach spaces and let W be a Banach space containing \mathcal{E} uniformly complemented. If Y is a Banach space complemented in its bidual such that $Ext(W, Y) = 0$, then $Ext(Z, Y) = 0$ for every Banach space Z locally \mathcal{E} .*

Jebreen et al proved the corresponding version for the colocality of Banach spaces as follows

Theorem 2 [18, Theorem 1.7] *Let \mathcal{E} be a family of finite dimensional Banach spaces and let W be a Banach space containing \mathcal{E} uniformly complemented. If Y is a Banach space such that $\text{Ext}(Y, W) = 0$, then $\text{Ext}(Y, Z) = 0$ for every Banach space Z complemented in its bidual and colocally \mathcal{E} .*

The triviality of all twisted sums of two Banach spaces is inherited by their complemented subspaces.

Proposition 3 [5, Lemma 3], [17, Proposition 2.3] *Let X, A_1 and A_2 be Banach spaces such that $X = A_1 \oplus A_2$. Then for any Banach space U*

- (i) *$\text{Ext}(U, X) = 0$ if and only if $\text{Ext}(U, A_i) = 0$ for $i = 1, 2$.*
- (ii) *$\text{Ext}(X, U) = 0$ if and only if $\text{Ext}(A_i, U) = 0$ for $i = 1, 2$.*

Throughout this paper \mathbb{K} denotes the scalar field; either $\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$ and c_{00} denotes the vector space of all finitely supported sequences in \mathbb{K} , that is $c_{00} := \{(\alpha_n) : \alpha_n \in \mathbb{K}, n \in \mathbb{N} \text{ and } \exists N \in \mathbb{N} : n = 0, \forall n > N\}$.

2. Twisted sums with James spaces.

In 1951, Robert C. James provided the first example of a non-reflexive Banach space isomorphic to its second dual, called the James space J_2 [15]. Edelstein and Mityagin were the first to observe that it can be generalized to an arbitrary $p > 1$ as they defined

$$\|x\|_{J_p} = \sup \left\{ \left(\sum_{i=1}^k |a_{n_i} - a_{n_{i+1}}|^p \right)^{\frac{1}{p}} : k, n_1, n_2, \dots, n_{k+1} \in \mathbb{N}, n_1 < n_2 < \dots < n_{k+1} \right\}$$

and the Banach space $\{x = (x_n)_{n \in \mathbb{N}} \in c_0 : \|x\|_{J_p} < \infty\}$ is called the p^{th} James Space J_p . It can be seen that the completion of c_{00} with respect to this norm is J_p [12]. Moreover, Edelstein and Mityagin showed that James' proof of the quasi-reflexivity of J_2 , the original James space, can be carried out to J_p for every $p > 1$ [12]. The proof can not, however, work for $p = 1$ because J_1 is isometrically isomorphic to ℓ_1 . Bird et al [4, 2.3] proved that J_p contains a complemented copy of ℓ_p , for $1 < p < \infty$.

Proposition 4 (i) *$\text{Ext}(J_p, \ell_1) \neq 0$, that is $\text{Ext}(J_p, J_1) \neq 0, 1 < p < \infty$.*
(ii) *$\text{Ext}(J_p, J_q) \neq 0, 1 < p, q < \infty$.*

Proof. (i) Note that c_0 is finitely represented in J_p [3, Theorem 1.1], that is for each $\epsilon > 0$, and each finite-dimensional subspace E of c_0 , there exists a subspace F of J_p , depending on E such that there is an isomorphism T on E onto F satisfying $\|T\| \|T^{-1}\| < 1 + \epsilon$ [3]. Hence J_p contains $\{\ell_\infty^n\}_{n=1}^\infty$ uniformly complemented. Since $\text{Ext}(c_0, \ell_1) \neq 0$ [6, Theorem 5.1], then the result can be deduced by Theorem 1.

(ii) For $1 < p < \infty$, J_p contains a complemented copy of ℓ_p [4,2.3], and ℓ_p contains $\{l_p^n\}$ uniformly complemented [23, II.5.9], then J_p contains $\{l_p^n\}$ uniformly complemented. Hence $\text{Ext}(\ell_p, \ell_q) \neq 0$, where $1 < p, q < \infty$ [7, Section 5], implies that $\text{Ext}(J_p, \ell_q) \neq 0$ by Theorem 1 and $\text{Ext}(\ell_p, J_q) \neq 0$ by Theorem 2. Therefore $\text{Ext}(J_p, J_q) \neq 0$ by Proposition 3. ■

The Schreier space S_1 was first considered by Schreier in 1930 [26], in order to provide an example of a weakly null sequence without Cesaro summable subsequence. A variation of this idea gave rise to the construction of the Schreier spaces [2], [8]. Bird and Laustsen generalized the concept of a Schreier space from one Schreier space, corresponding to the ℓ_1 -norm, to a whole family, one for each $p \geq 2$, corresponding to the ℓ_p -norms as follows:

$$\|x\|_{Z_p} = \sup_A \left(\sum_{j \in A} |x_j|^p \right)^{\frac{1}{p}}$$

where $x = (x_n)_{n \in \mathbb{N}}$ and the supremum is taken over all admissible subsets of \mathbb{N} , which are defined as the finite subsets $A = \{n_1, n_2, \dots, n_k\}$ of N such that $k \leq n_1 < n_2 < \dots < n_k$. The subspace $\{x = (x_n)_{n \in \mathbb{N}} \in c_0 : \|x\|_{Z_p} < \infty\}$ of $\mathbb{K}^{\mathbb{N}}$ is a Banach space called the unrestricted p^{th} Schreier Space Z_p . The completion of c_{00} with respect to $\|x\|_{Z_p}$ is the restricted p^{th} Schreier Space S_p [4,3.2,3.6].

In 2010, Bird and Laustsen create a new family of Banach spaces, the James-Schreier spaces, by amalgamating the two important classical Banach spaces: James' quasi-reflexive Banach space and Schreier's Banach space and they proved that these spaces are counterexamples to the Banach-Saks property and that most of the results about the James space as a Banach algebra carry over to the new spaces; see [4] for details. For $1 \leq p < \infty$, they defined the following norm:

$$\|x\|_{W_p} = \sup_A \left(\sum_{i=1}^k |x_{n_i} - x_{n_{i+1}}|^p \right)^{\frac{1}{p}}$$

where the supremum is taken over all permissible subsets of \mathbb{N} , which are defined as the finite subsets $A = \{n_1, n_2, \dots, n_{k+1}\}$ of N such that $k \leq n_1 < n_2 < \dots < n_{k+1}$. The subspace $Z_p = \{x = (x_n)_{n \in \mathbb{N}} \in c_0 : \|x\|_{W_p} < \infty\}$ of $\mathbb{K}^{\mathbb{N}}$ is a Banach space called the unrestricted p^{th} James-Schreier Space W_p . The completion of c_{00} with respect to $\|x\|_{W_p}$ is the restricted p^{th} James-Schreier Space V_p [4,4.2,4.8].

Proposition 5 Let $U \in \{\ell_q, S_q, V_q, Z_q, W_q, V_q^{**}, 1 < q < \infty\}$. Then

- (i) $\text{Ext}(J_p, U) \neq 0, 1 < p < \infty$.
- (ii) $\text{Ext}(U, J_p) \neq 0, 1 < p < \infty$.

Proof. Since $\text{Ext}(\ell_p, U) \neq 0$, and $\text{Ext}(U, \ell_p) \neq 0, 1 < p < \infty$, where U is either $\ell_q, S_q, V_q, Z_q, W_q$, or V_q^{**} by [19, Proposition 2.4], and ℓ_p is complemented in J_p

then $Ext(J_p, U) \neq 0$ and $Ext(U, J_p) \neq 0$ by Proposition 3. ■

The original James space J_2 has an additional twisted sums due to $Ext(U, \ell_2) \neq 0$, where U is either c_0, S_1, V_1, Z_1, W_1 or V_1^{**} [19, Proposition 2.3] which gives

Proposition 6 $Ext(U, J_2) \neq 0$ where $U \in \{c_0, S_1, V_1, Z_1, W_1, V_1^{**}\}$.

3. Twisted sums of the Tsirelson's space and the AD space.

A Banach space X is said to be asymptotic ℓ_p , $1 \leq p \leq \infty$ with respect to a basis $(e_i)_{i=1}^\infty$, if there exists a constant $C \geq 1$ (the asymptotic constant), such that $(x_i)_{i=1}^n$ is C -equivalent to the unit vector basis of ℓ_p^n , i.e. for any n -tuple of scalars $a = (a_1, \dots, a_n)$,

$$\frac{1}{C} \|a\|_p \leq \left\| \sum_{i=1}^n a_i x_i \right\| \leq C \|a\|_p$$

In 1974, Tsirelson [27] constructed the first example of a reflexive Banach space with unconditional basis that is asymptotically ℓ_∞ but has no embedded copy of c_0 or ℓ_p ($1 \leq p < \infty$). He constructed a convex, weakly compact subset V of c_0 with the following properties:

- (i) For all $n \in \mathbb{N}, e_n \in V$,
- (ii) If $f = (f_n) \in V$ and $g = (g_n)$ is such that $|g_n| \leq |f_n|$ for all $n \in \mathbb{N}$, then $g \in V$,
- (iii) If $f_1, \dots, f_n \in V$ are such that $n \leq f_1 < \dots < f_n$, then $\frac{1}{2}(f_1 + \dots + f_n) \in V$,
- (iv) For all $x \in V$ there exists $n \in \mathbb{N}$ such that $2P_n(x) \in V$.

Tsirelson's space is the linear span of the set V with the norm that makes V be the unit ball. Figiel and Johnson [14] constructed the conjugate of Tsirelson example, known as the T space. It is defined as the closure of the finitely supported sequences with the norm $\|x\|_T := \lim \|x\|_m$, where the norms $\|x\|_m$ are defined inductively:

$$\|x\|_0 = \|x\|_{c_0}, \|x\|_{m+1} = \left\{ \max \|x\|_m, \frac{1}{2} \max \left[\sum_{j=1}^n \|E_j x\|_m \right] \right\}$$

and the inner max is taken over all choices of consecutive finite sets $\{E_j\}_{j=1}^n$, $n \leq E_1 < E_2 < \dots < E_n$ [14]. The original Tsirelson space is denoted in the literature by T^* .

Recall that $\{I_\infty^n\} \subseteq c_0$, for each $n \in \mathbb{N}$, hence c_0 is locally $\{I_\infty^n\}$.

Proposition 7 $Ext(T^*, U) \neq 0$, where $U \in \{\ell_q, S_q, Z_q, W_q, V_q, V_q^{**}, J_q, q = 1, 2\}$.

Proof. Since T^* is asymptotically ℓ_∞ then T^* contains $\{I_\infty^n\}$ uniformly complemented. Hence $Ext(c_0, \ell_1) \neq 0$ [7, Theorem 5.1], and $Ext(c_0, \ell_2) \neq 0$ [6, 4.2] imply $Ext(T^*, \ell_1) \neq 0$, and $Ext(T^*, \ell_2) \neq 0$ by Theorem 1. For $U \in \{\ell_q, S_q, Z_q, W_q, V_q, V_q^{**}, J_q, q = 1, 2\}$, U contains $\{\ell_q^n\}$ uniformly complemented Theorem 2. ■

Argyros and Deliyanni constructed two examples of asymptotic ℓ_1 Banach spaces, both are of the type of Tsirelson's [1]. The second, we shall denote it by AD , does not contain any unconditional basic sequence.

Proposition 8 (i) $Ext(U, T) \neq 0$, and $Ext(U, AD) \neq 0$ where $U \in \{c_0, S_p, V_p, Z_p, W_p, V_p^{**}, 1 \leq p < \infty\}$.
(ii) $Ext(T^*, T) \neq 0$, and $Ext(T^*, AD) \neq 0$

Proof. (i) By [19, Proposition 2.2], $Ext(U, \ell_1) \neq 0$ where $U \in \{c_0, S_p, V_p, Z_p, W_p, V_p^{**}, 1 \leq p < \infty\}$. Since T and AD are asymptotically ℓ_1 , then T and AD contain $\{\ell_1^n\}$ uniformly complemented. The result follows by Theorem 2.

(ii) Since $Ext(T^*, \ell_1) \neq 0$ by the previous proposition, and since the spaces T and AD contain $\{\ell_1^n\}$ uniformly complemented, the result follows by Theorem 2. ■

4. Twisted sums of the Johnson-Lindenstrauss space.

Johnson and Lindenstrauss [20] constructed a nontrivial twisted sum of c_0 and a Hilbert space (necessarily non-separable), called the Johnson-Lindenstrauss space JL . It is defined to be the completion of the linear span of $c_0 \cup \{\chi_i : i \in I\}$ in ℓ_∞ with respect to the norm:

$$\|y = x + \sum_{j=1}^k a_{i(j)} \chi_{i(j)}\| = \max \left\{ \|y\|_\infty, \|(a_i)_{i \in I}\|_{\ell_2(I)} \right\}$$

$x \in c_0$, $a_{i(j)}$ are scalars, and χ_i is the characteristic function of A_i , $\{A_i\}_{i \in I}$ is an almost disjoint uncountable family of infinite subsets of \mathbb{N} . They proved that JL/c_0 is isomorphic to some $\ell_2(I)$ and since ℓ_1 is projective, the dual sequence $0 \rightarrow \ell_2(I) \rightarrow JL^* \rightarrow \ell_1 \rightarrow 0$ of the exact sequence $0 \rightarrow c_0 \rightarrow JL \rightarrow \ell_2(I) \rightarrow 0$ splits. That is, $JL^* = \ell_1 \oplus \ell_2(I)$.

Proposition 9 (i) $Ext(JL^*, U) \neq 0$, where $U \in \{\ell_1, S_1, Z_1, W_1, V_1, V_1^{**}, T, AD\}$.
(ii) $Ext(U, JL^*) \neq 0$, where $U \in \{T^*, c_0, S_p, Z_p, W_p, V_p, V_p^{**}, 1 \leq p < \infty\}$.

Proof. (i) Since $JL^* = \ell_1 \oplus \ell_2(I)$ and ℓ_2 is complemented in $\ell_2(I)$, then ℓ_2 is complemented in JL^* . By [6, 4.3] $Ext(\ell_2, \ell_1) \neq 0$, hence $Ext(JL^*, \ell_1) \neq 0$, which leads to $Ext(JL^*, T) \neq 0$, $Ext(JL^*, AD) \neq 0$, $Ext(JL^*, S_1) \neq 0$ and $Ext(JL^*, Z_1) \neq 0$ by Theorem 2. By proposition 3, the result follows.

(ii) Since ℓ_1 is complemented in JL^* , then the result can be concluded using $Ext(c_0, \ell_1) \neq 0$ and a similar argument to that used in (i). ■

In [17, Theorem 2.2] it has been proved that for any Banach spaces U and Y , $Ext(Y, U^*) = 0$ if and only if $Ext(U, Y^*) = 0$. Therefore it is immediate that

Corollary 10 $Ext(JL, U^*) \neq 0$, where $U \in \{T^*, c_0, S_p, Z_p, W_p, V_p, V_p^{**}, 1 \leq p < \infty\}$.

5. Twisted sums of the James tree space.

The James tree space JT was introduced by James in [15]. It was the first example of a separable dual Banach space that contains no copy of ℓ_1 though it has a non separable dual. Moreover, James proved that JT does not contain

a subspace isomorphic to c_0 or ℓ_1 . The space JT is defined to be the completion of the space of finite sequences over the dyadic tree Δ with respect to the norm

$$\|x\| = \sup_{n \in \mathbb{N}} \sup_{S_1, \dots, S_n} \left[\sum_{i=1}^n \left(\sum_{\alpha \in S_i} x_\alpha \right)^2 \right]^{1/2}$$

where the supremum is taken over all finite sets of pair wise disjoint segments of Δ .

Proposition 11 (i) $Ext(JT, U) \neq 0$, where $U \in \{\ell_1, S_1, Z_1, W_1, V_1, V_1^{**}, T, AD, JL^*\}$.

(ii) $Ext(JT, U) \neq 0$, where $U \in \{\ell_2, S_2, Z_2, W_2, V_2, V_2^{**}, JL^*\}$.

Proof. (i) Fetter de Buen [13, 2.b.8, 3.a.7] proved that c_0 is finitely represented in the James space J . Since JT contains J , then JT contains $\{\ell_\infty^n\}_{n=1}^\infty$ uniformly which implies that it contains $\{\ell_\infty^n\}_{n=1}^\infty$ uniformly complemented. Since $Ext(c_0, \ell_1) \neq 0$, then by Theorem 1 we have $Ext(JT, \ell_1) \neq 0$. Applying Theorem 2 gives $Ext(JT, T) \neq 0, Ext(JT, AD) \neq 0, Ext(JT, S_1) \neq 0$ and $Ext(JT, Z_1) \neq 0$. The result follows by Proposition 3.

(ii) The proof is by using $Ext(c_0, \ell_2) \neq 0$ and applying Theorems 1, 2, and Proposition 3. ■

Proposition 12 $Ext(U, \mathcal{B}) \neq 0$, where \mathcal{B} is the predual of JT , and $Ext(U, JT^*) \neq 0$, where $U \in \{\ell_p, S_p, Z_p, W_p, V_p, V_p^{**}, 1 \leq p < \infty\}$.

Proof. The predual B of JT contains $\{\ell_1^n\}_{n=1}^\infty$ uniformly complemented [18, Lemma 2.4], and hence so does JT^* . It has been proved in [19, Proposition 2.2] that $Ext(U, \ell_1) \neq 0$, where U is either $\ell_p, S_p, V_p, Z_p, W_p$, or V_p^{**} , so by Theorem 2 we get the result. ■

6. Singular Twisted sums with $C(K)$ spaces.

A continuous linear operator $T : E \rightarrow F$ between two Banach spaces is called strictly singular if it fails to be invertible on any infinite dimensional closed subspace of E . We say that a quasi-linear map $F : U \rightarrow Y$ is strictly singular if it has no trivial restriction to any infinite dimensional subspace of U . A quasi-linear map $F : U \rightarrow Y$ is strictly singular if and only if the quotient map $Q : Y \oplus_F U \rightarrow U$ is a strictly singular operator, that is the restriction of Q to any infinite dimensional subspace of $Y \oplus_F U$ is not an isomorphism [11, Lemma 1]. In this case we say that the twisted sum $Y \oplus_F U$ is singular.

Theorem 13 There are singular twisted sums $U \oplus_F C[0, 1]$, where $U \in \{T, T^*, AD, JT\}$.

Proof. For every $U \in \{T, T^*, AD, JT\}$, U is separable and have no copy of ℓ_1 , hence there is an exact sequence

$$0 \rightarrow C[0, 1] \xrightarrow{j} X \xrightarrow{Q} U \rightarrow 0$$

with Q strictly singular [7, 2.3]. Hence the corresponding quasi-linear maps $F : U \rightarrow C[0, 1]$ is strictly singular. ■

Recall that if $N \in \mathbb{N}$, the space $C(\omega^N)$ is isomorphic to c_0 , and so, by Sobczyk's Theorem, for any separable Banach space U , we have $\text{Ext}(U, C(\omega^N)) = 0$. The extension constant $\pi_N(U)$ is the least constant such that if

$$0 \rightarrow C(\omega^N) \xrightarrow{J} X \xrightarrow{Q} U \rightarrow 0$$

is an exact sequence and $\varepsilon > 0$, then there is a linear operator $P : X \rightarrow C(\omega^N)$ with $PJ = i_{C(\omega^N)}$ and $\|P\| \leq \pi_N(U) + \varepsilon$ [7, Section 3]. It is proved that $\pi_N(U) \leq 2N + 1$, for every $N \in \mathbb{N}$ [7, Theorem 3.1].

Theorem 14 $\text{Ext}(T, C(\omega^\omega)) = 0$.

Proof. Recall that T^* is a separable Banach space with summable Szlenk index [21, Proposition 6.7]. Hence $\text{Ext}(T, C(\omega^\omega)) = 0$ by [7, 4.4], which implies that $\text{Sup}_N \pi_N(U) < \infty$ by [7, 4.1]. ■

7. Pełczyński's property (u).

Two infinite-dimensional Banach spaces X and Y are totally incomparable if no closed, infinite-dimensional subspace of X is isomorphic to a subspace of Y . Since $\{J_p, S_q\}$ and $\{J_p, V_q\}$ are totally incomparable for $p, q \geq 1$ [3, 5.9]; we can deduce immediately that any twisted sum that extends J_p can not be isomorphic to S_q or V_q , $p, q \geq 1$. But we can know more about the twisted sums of certain spaces using Pełczyński's property (u). A Banach space X has Pełczyński's property (u) if for every weak Cauchy sequence $(x_n)_{n \in \mathbb{N}}$ in X , there is a sequence $(y_n)_{n \in \mathbb{N}}$ in X such that for every bounded functional on X we have

$$\sum_{n=1}^{\infty} |\langle y_n, f \rangle| < \infty \text{ and } \left\langle x_n - \sum_{i=1}^n y_i, f \right\rangle \rightarrow 0 \text{ as } n \rightarrow \infty$$

We will show that for any Banach space U , every twisted sum that extends J_2 or the p^{th} -James-Schreier space V_p ($p > 1$) does not have the Pełczyński's property (u). For this purpose we need the Pełczyński's Theorem:

Theorem 15 (22) (i) Every Banach space with an unconditional basis has Pełczyński's property (u).

(ii) Every closed subspace of a Banach space with Pełczyński's property (u) has Pełczyński's property (u).

(iii) The James space J_2 does not have Pełczyński's property (u).

Theorem 16 The twisted sum that extends V_p ($p > 1$) or J_2 does not have the Pełczyński's property (u), and hence has no unconditional basis.

Proof. Let $X \in \text{Ext}(U, V_p)$, then V_p is isomorphic to a closed subspace of X . Since V_p does not have the Pełczyński's property (u) [4, Theorem 6.3] then X does not have Pełczyński's property (u), by (ii) of Pełczyński theorem. Hence X has no unconditional basis. The case for $X \in \text{Ext}(U, j_2)$ can be proved similarly by using (iii) of the previous theorem. ■

Data Availability

The readers who are interested may contact the author for more informations.

Conflicts of Interest

The author declares no conflict of interest.

References

- [1] S.A. Argyros and I. Deliyanni, Examples of Asymptotic ℓ_1 Banach spaces, *Journal of Functional Analysis* 159(1)(1997), 43-109.
- [2] B. Beauzamy and J. T. Lapreste, Modèles étalés des espaces de Banach, *Publications du Département de mathématiques (Lyon)*, no. 4A (1983) 1-199, .
- [3] A. Bird, G. Jameson and N. J. Laustsen, The Giesy-James theorem for general index p , with an application to operator ideals on the p^{th} James space, *J. Operator Theory* 70 (2013), 291–307.
- [4] A. Bird and N. Laustsen, An amalgamation of the Banach spaces associated with James and Schreier, Part I : Banach-space structure, *Banach Center Publications*, 91 (2010) 45-76.
- [5] F. Cabello Sanchez and J.M. Castillo, Duality and twisted sums of Banach spaces, *J.Funct. Anal.* 175 (2000) 1-16.
- [6] F. Cabello Sanchez and J. M. Castillo, Uniform boundedness and twisted sums of Banach spaces, *Houston J. Math.* 30 (2) (2004) 523-536.
- [7] F. Cabello Sanchez, J. M. Castillo, N. J. Kalton and D. T. Yost, Twisted sums with $C(K)$ spaces, *Trans. Amer. Math. Soc.* 355 (11) (2003) 4523—4541.
- [8] P. G. Casazza and T. J. Shura, Tsirelson's space, *Lecture Notes in Mathematics*, vol. 1363 (1989), Springer-Verlag, Berlin.
- [9] J. M. Castillo, Nonseparable $C(K)$ -spaces can be twisted when K is a finite height compact, *Topology and its Applications* 198 (2016) 107–116.
- [10] J. M. Castillo and M. Gonzalez, *Three-space Problems in Banach Space Theory*, Springer Lecture Notes in Mathematics 1667(1997).

- [11] J. M. Castillo and Y. M. Salguero, Strictly singular quasi-linear maps, *Non-linear Analysis* 49 (2001) 897 – 903.
- [12] Edelstein and Mityagin, Homotopy type of linear groups of two classes of Banach spaces, *Functional Anal. Appl.* 4 (1970), 221–231.
- [13] H. Fetter, B. Gamboa de Buen, *The James forest*, London Mathematical Society Lecture Note Series. 236. Cambridge: Cambridge University Press (1997).
- [14] T. Figiel and W. B. Johnson, A uniformly convex Banach space which contains no ℓ_p . *Compositio Mathematica*, Volume 29 (1974) no. 2, pp. 179–190.
- [15] R. C. James, A non-reflexive Banach space isometric with its second conjugate space. *Proc. Nat. Acad. Sci. U. S. A.* 37, (1951). 174–177.
- [16] R. C. James, A separable somewhat reflexive Banach space with nonseparable dual, *Bull. Amer. Math. Soc.* 80 (1974), 738–743.
- [17] F. Jamjoom and H. Jebreen, On Twisted Sums Of Banach Spaces, *Far East Journal of Mathematical Sciences (FJMS)*, 26(1) (2007)169-184.
- [18] H. Jebreen, F. Jamjoom and D. Yost, Colocality and Twisted Sums, *J. Math. Anal. & App.* 323 (2) (2006) 864-875.
- [19] H. Jebreen, *Proceedings of the 7th International Conference on Applied Analysis and Mathematical Modeling, ICAAMM18, Istanbul-Turkey.*
- [20] W.B. Johnson, and J. Lindenstrauss, Some remarks on weakly compactly generated Banach spaces *Israel J. Math.* 17 (1974), 219–230.
- [21] H. Knaust, E. Odell and T. Schlumprecht, On asymptotic structure, the Szlenk index and UKK properties in Banach spaces, *Positivity* 3 (1999), no. 1, 173-199.
- [22] J. Lindenstrauss and C. Stegall: Examples of separable spaces which do not contain 1 and whose duals are nonseparable, *Studia Math.* 54, (1975) 81-105.
- [23] J. Lindenstrauss and L. Tzafriri, Classical Banach spaces. *Bull. Amer. Math. Soc.* 81 (1975), no. 1.
- [24] R. E. Megginson, *An introduction to Banach space theory*, Graduate Texts in Mathematics Vol. 183 (1998), Springer Verlag.
- [25] A. Pełczyński, A connection between weakly unconditional convergence and weakly completeness of Banach spaces, *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astr. Phys.* 6 (1958), 251–253.

- [26] J. Schreier, Ein Gegenbeispiel zur theorie der schwachen konvergenz, *Studia Math.*, 2, (1930), 58–62.
- [27] B.S. Tsirelson, Not every Banach space contains an imbedding of ℓ_p , or c_0 , *Functional Analysis and its Applications* 8 (1974), no. 2, 138-141.